

On the United Theory of the Family of Euler-Halley Type Methods with Cubical Convergence in Banach Spaces*¹⁾

Xinghua Wang

(Department of Mathematics, Zhejiang University, Hangzhou 310028, China)
(Wenzhou University, Wenzhou 325027, P. R. China)

Chong Li

(Department of Mathematics, Zhejiang University Hangzhou 310027, China)
(Academy of Mathematics and System Sciences, CAS)

Abstract

The convergence problem of the family of Euler-Halley methods is considered under the Lipschitz condition with the L -average, and a united convergence theory with its applications is presented.

Key words: Operator equation, The family of Euler-Halley, Iterations, Cubical convergence

1. Introduction

Let E and F be real or complex Banach space with norm $\|\cdot\|$, and let $f : D \subset E \rightarrow F$ be a nonlinear twice differentiable operator. The family of Euler-Halley iterations with the parameter $\lambda \in [0, 2]$ for solving the operator equation $f(x) = 0$ is defined as follows:

$$x_{n+1} = T_{f,\lambda}(x_n) = x_n + u_f(x_n) + v_{f,\lambda}(x_n), \quad n = 0, 1, \dots, \quad (1.1)$$

where

$$\begin{aligned} u_f(x) &= -f'(x)^{-1}f(x), \\ v_{f,\lambda}(x) &= -\frac{1}{2}f'(x)^{-1}f''(x)u_f(x)Q_{f,\lambda}(x)u_f(x), \\ Q_{f,\lambda}(x) &= \left\{I + \frac{\lambda}{2}f'(x)^{-1}f''(x)u_f(x)\right\}^{-1}. \end{aligned}$$

This family includes, as particular cases, the well known Euler method ($\lambda = 0$), [1, 4, 12], the Halley method ($\lambda = 1$), [3, 5, 10, 12, 18] and the convex acceleration of Newton's method or super-Halley method ($\lambda = 2$), [6, 7, 11], so that recent interests are focused in this direction, see for example [2, 8, 9]. In particular, using a quadratic majorizing function, Argyros et al analyze the convergence of the method (1.1). However it is incorrect as shown by Han [9]. In [8], Gutierrez and Hernandez established the convergence with a cubic polynomial as the majorizing function under the classical Lipschitz condition of f'' while Han [9] established the convergence under the weak condition, so-called, γ -condition of f'' , which was first presented by Wang [13, 14] when he investigated the convergence of the family of Halley methods. The purpose of the present paper is to give a united convergence theory for the family of Euler-Halley iterations such that all the known results are included as its special cases. Also some new results are obtained as the corollaries. It should be noted that this work is in spirit of Wang's idea in [15, 16].

2. Preliminaries and Lemmas

Let $D \subset E$ be a convex subset, open or closed. For $x_0 \in E, r > 0$, let $B(x_0, r)$ denote the open ball with the radius r and the center x_0 while the corresponding closed ball is denoted

* Received May 17, 2000.

¹⁾This project is supported by the Special Funds for Major State Basic Research Projects (Grant No. G19990328) and the National Natural Science Foundation of China (Grant No. 10271025); also supported partly by Zhejiang Provincial Natural Science Foundation of China.

by $\overline{B(x_0, r)}$. Through the paper, we always assume that $f'(x_0)^{-1}$ exists. In order to study the convergence we require some definitions and lemmas, some of which are directly taken from [15, 16].

Definition 2.1^[15]. A function f from D to F is called to satisfy the center Lipschitz condition in the ball $B(x_0, r)$ with the L average if

$$\|f(x) - f(x_0)\| \leq \int_0^{\rho(x)} L(u)du, \quad \forall x \in B(x_0, r), \quad (2.1)$$

where $\rho(x) = \|x - x_0\|$ and L is a positive integrable function on the interval $[0, R]$ for some sufficient large number $R > 0$, for example, with $\int_0^R (R - u)L(u)du = R$.

Take $r_0 > 0$ such that

$$\int_0^{r_0} L(u)du = 1 \quad (2.2)$$

and set

$$b = \int_0^{r_0} uL(u)du. \quad (2.3)$$

For $\beta \in (0, b]$, define

$$h(t) = \beta - t + \int_0^t (t - u)L(u)du, \quad \forall t \in [0, R]. \quad (2.4)$$

Lemma 2.1^[15]. The function h is decreasing monotonically in $[0, r_0]$, while it is increasing monotonically in $[r_0, R]$. Moreover, if $\beta \leq b$,

$$h(\beta) > 0, \quad h(r_0) = \beta - b \leq 0, \quad h(R) = \beta > 0.$$

Consequently, h has a unique zero in each interval, respectively, which are denoted by r_1 and r_2 . They satisfy

$$\beta < r_1 < \frac{r_0}{b}\beta < r_0 < r_2 < R \quad (2.5)$$

when $\beta < b$ and $r_1 = r_2$ when $\beta = b$.

Furthermore, we assume that L is a positive nondecreasing differentiable function in $[0, R]$. Then we have the following lemma.

Lemma 2.2. Let h be defined as (2.4) and $\beta \leq b$. Then, for each $t \in [0, r_1]$,

(i) $H_h(t) = h'(t)^{-2}h''(t)h(t) < 1$;

(ii) $T_{h,\lambda}(t) \in [0, r_1]$;

(iii) $t \leq T_{h,\lambda}(t)$.

Proof. (i) It suffices to show that

$$g(t) = h'(t)^2 - h''(t)h(t) > 0.$$

Since

$$g'(t) = h'(t)h''(t) - h'''(t)h(t) = h'(t)L(t) - L'(t)h(t) \leq 0,$$

so that $g(t) \geq g(r_1) = h'(r_1)^2 > 0$ and proves (i).

(ii) Observe that

$$T'_{h,\lambda}(t) = \frac{H_h(t)^2}{2(1 - \frac{\lambda}{2}H_h(t))^2} [3(1 - \frac{\lambda}{2}) + \frac{\lambda}{2}(\lambda - 1)H_h(t) - H_{h'}(t)].$$

Since $H_{h'}(t)$ is negative and $0 < H_h(t) < 1$ for each $t \in [0, r_1]$, it follows that $T'_{h,\lambda}(t) > 0$ for all $t \in [0, r_1]$ and each $\lambda \in [1, 2]$. Hence $T_{h,\lambda}(t)$ is monotonically increasing on $[0, r_1]$ for each $\lambda \in [1, 2]$. Consequently, $T_{h,\lambda}(t) \leq T_{h,\lambda}(r_1) = r_1$ for each $\lambda \in [1, 2]$. On the other hand, for any $\lambda \in [0, 1]$, we have

$$T_{h,\lambda}(t) \leq T_{h,1}(t) \leq r_1.$$

Thus (ii) holds.

(iii) This results from that $u_h(t) \geq 0$ and $v_{h,\lambda}(t) \geq 0$. ■

The following two lemmas are due to [16]. Note that, under the assumption of the following Lemma 2.3, we may assume that $f'(x_0)^{-1}f'$ satisfies the center Lipschitz condition in the closed ball $\overline{B(x_0, r)}$ with the L average.

Lemma 2.3. Suppose that $f'(x_0)^{-1}f'$ satisfies the center Lipschitz condition in the ball $B(x_0, r)$ with the L average and $r > r_0$. Then for each $x \in B(x_0, r_0)$, $f'(x)^{-1}$ exists and

$$\|f'(x)^{-1}f'(x_0)\| \leq \frac{1}{1 - \int_0^{\rho(x)} L(u)du}. \quad (2.6)$$

Lemma 2.4. Let $\beta = \|f'(x_0)^{-1}f(x_0)\| \leq b$. Assume that $r_1 \leq r < r_2$ if $\beta < b$, or $r = r_1$ if $\beta = b$. Then, under the hypotheses of the conditions (2.1), the equation $f(x) = 0$ has a unique solution

$$x^* \in \overline{B(x_0 - f'(x_0)^{-1}f(x_0), r_1 - \beta)} \subset \overline{B(x_0, r_1)} \quad (2.7)$$

in the closed ball $\overline{B(x_0, r)}$.

We still need a lemma below.

Lemma 2.5. Suppose that

$$\|f'(x_0)^{-1}f''(x_0)\| = L(0) \quad (2.8)$$

and

$$\|f'(x_0)^{-1}(f''(x') - f''(x))\| \leq L(\rho(\overline{xx'})) - L(\rho(x)), \quad (2.9)$$

$$\forall x \in \overline{B(x_0, r)}, \quad \forall x' \in \overline{B(x, r - \rho(x))},$$

where $\rho(x) = \|x - x_0\|$, $\rho(\overline{xx'}) = \rho(x) + \|x' - x\|$. Then, for each $x \in B(x_0, r_0)$,

(i) $\|f'(x_0)^{-1}f''(x)\| \leq h''(\|x - x_0\|)$;

(ii) $f'(x)^{-1}$ exists and

$$\|f'(x)^{-1}f'(x_0)\| \leq \frac{1}{h'(\|x - x_0\|)}.$$

Proof. It follows from (2.8) and (2.9) that

$$\|f'(x_0)^{-1}f''(x)\| \leq L(0) + \|f'(x_0)^{-1}(f''(x) - f''(x_0))\| \leq L(\|x - x_0\|) = h''(\|x - x_0\|),$$

using the fact that $h''(t) = L(t)$ and (i) follows.

To prove (ii), by Lemma 2.2, it suffices to show that $f'(x_0)^{-1}f'$ satisfies the center Lipschitz condition in the closed ball $B(x_0, r_0)$ with the L average since $h'(\|x - x_0\|) = -1 + \int_0^{\|x - x_0\|} L(u)du$. By Taylor formula,

$$f'(x) = f'(x_0) + f''(x_0)(x - x_0) + \int_0^1 (f''(x_0 + t(x - x_0)) - f''(x_0)) dt(x - x_0).$$

This, together with (2.8) and (2.9), implies that

$$\begin{aligned} \|f'(x_0)^{-1}f'(x) - I\| &\leq L(0)\|x - x_0\| + \int_0^1 \|f''(x_0 + t(x - x_0)) - f''(x_0)\| dt\|x - x_0\| \\ &\leq \int_0^{\rho(x)} L(u)du. \end{aligned}$$

Hence (ii) holds and the proof is complete. ■

3. The United Convergence Theorem

Let $\{x_n\}$ be defined as (1.1) and let $\{t_n\}$ be the corresponding sequence of the majorizing function h , that is,

$$t_{n+1} = T_{h,\lambda}(t_n), \quad n = 0, 1, \dots, \quad (3.1)$$

where $t_0 = 0$. Then it follows from Lemma 2.2 that $\{t_n\}$ is well defined, increasing monotonically and tending to r_1 .

Lemma 3.1^[8]. For any $n = 0, 1, 2, \dots$,

$$f(x_{n+1}) = \frac{1}{2}f''(x_n)\{(2-\lambda)u_f(x_n) + v_{f,\lambda}(x_n)\}v_{f,\lambda}(x_n) + \int_0^1 \{f''(x_n + \tau(x_{n+1} - x_n)) - f''(x_n)\}(1-\tau)d\tau(x_{n+1} - x_n)^2. \quad (3.2)$$

Now we are ready to give the main theorem of this section.

Theorem 3.1. Suppose that $r > r_1$ and (2.8),(2.9) hold on $\overline{B(x_0, r)}$. If

$$\beta = \|f'(x_0)^{-1}f(x_0)\| \leq b = \int_0^{r_0} uL(u)du,$$

then the iteration

$$x_{n+1} = T_{f,\lambda}(x_n), \quad n = 0, 1, \dots,$$

is well defined for all $\lambda \in [0, 2]$ and that $\{x_n\}$ converges to a solution x^* of the equation $f(x) = 0$ satisfying

$$x^* \in \overline{B(x_0 - f'(x_0)^{-1}f(x_0), r_1 - \beta)} \subset \overline{B(x_0, r_1)}. \quad (3.8)$$

Moreover, for each r satisfying $r_1 \leq r < r_2$ if $\beta < b$ and $r = r_1$ if $\beta = b$, the equation $f(x) = 0$ has a unique solution in the closed ball $\overline{B(x_0, r)}$.

Proof. Using the mathematical induction, we can prove that following four statements hold for any $n = 0, 1, \dots$:

- (a) $f'(x_n)^{-1}$ exists and $\|f'(x_n)^{-1}f'(x_0)\| \leq h'(t_n)^{-1}h'(t_0)$;
- (b) $\|u_f(x_n)\| \leq u_h(t_n)$;
- (c) $Q_{f,\lambda}(x_n)$ exists and $\|Q_{f,\lambda}(x_n)\| \leq Q_{h,\lambda}(t_n)$;
- (d) $\|v_{f,\lambda}(x_n)\| \leq v_{h,\lambda}(t_n)$.

In fact, (a)-(b) are clearly true for $n = 0$. Consequently, $\|x_1 - x_0\| \leq t_1$. Now assume (a)-(b) are also true for n . Then

$$\|x_{n+1} - x_n\| \leq t_{n+1} - t_n \quad (3.9)$$

so that $\|x_{n+1} - x_0\| \leq t_{n+1} < r_1$. Thus Lemma 2.5 and Lemma 3.1 imply that (a) and (b) hold for $n + 1$. This again implies that (c) and (d) hold for $n + 1$. Hence x_{n+1} is well defined for all $n = 0, 1, \dots$ and $\lambda \in [0, 2]$. Furthermore, we also have that (3.9) holds for any $n = 0, 1, 2, \dots$ so that $\{x_n\}$ converges to a solution x^* of the equation $f(x) = 0$ satisfying $x^* \in \overline{B(x_0, r)}$ while (3.8) and the uniqueness of the solution x^* result from Lemma 2.4. The proof is complete. ■

4. Corollaries of the Convergence Theorem

In this section we will take L to be some particular functions and then obtain a series of concrete results, some of which have been given by some authors, see for example [8, 9], while the others seem new up to the present.

4.1 Kantorovith's Type Theorem

Given fixed positive constants γ and K , take

$$L(u) = \gamma + Ku. \quad (4.1)$$

Then r_0 is the solution of the equation

$$\int_0^{r_0} (\gamma + Ku)du = \gamma r_0 + \frac{1}{2}Kr_0^2 = 1,$$

i.e.,

$$r_0 = \frac{2}{\gamma + \sqrt{\gamma^2 + 2K}}. \quad (4.2)$$

Therefore

$$b = \int_0^{r_0} u(\gamma + Ku)du = \frac{2(\gamma + 2\sqrt{\gamma^2 + 2K})}{3(\gamma + \sqrt{\gamma^2 + 2K})^2}. \quad (4.3)$$

In this case the majorizing function is

$$h(t) = \beta - t + \frac{1}{2}\gamma t^2 + \frac{1}{6}Kt^3, \tag{4.4}$$

and $r_1 \leq r_2$ are its two positive solutions when $\beta \leq b$. Thus from Theorem 3.1 we immediately obtain the Kantorovich's type theorem.

Theorem 4.1. Suppose that $r > r_1$, $\|f'(x_0)^{-1}f''(x_0)\| = \gamma$ and

$$\begin{aligned} \|f'(x_0)^{-1}(f''(x') - f''(x))\| &\leq K\|x' - x\|, \\ \forall x \in \overline{B(x_0, r)}, \quad \forall x' \in \overline{B(x, r - \|x - x_0\|)}. \end{aligned} \tag{4.5}$$

If

$$\beta = \|f'(x_0)^{-1}f(x_0)\| \leq \frac{2(\gamma + 2\sqrt{\gamma^2 + 2K})}{3(\gamma + \sqrt{\gamma^2 + 2K})^2}, \tag{4.6}$$

then the iteration

$$x_{n+1} = T_{f,\lambda}(x_n), \quad n = 0, 1, \dots,$$

is well defined for all $\lambda \in [0, 2]$ and $\{x_n\}$ converges to a solution x^* of the equation $f(x) = 0$ satisfying

$$x^* \in \overline{B(x_0 - f'(x_0)^{-1}f(x_0), r_1 - \beta)} \subset \overline{B(x_0, r_1)}.$$

Moreover, for each r satisfying $r_1 \leq r < r_2$ if $\beta < b$ and $r = r_1$ if $\beta = b$, the equation $f(x) = 0$ has a unique solution in the closed ball $\overline{B(x_0, r)}$.

4.2 Smale's Type Theorem

For fixed $\gamma > 0$, let

$$L(u) = \frac{2\gamma}{(1 - \gamma u)^3}. \tag{4.7}$$

Then by [15] we have

$$r_0 = \left(1 - \frac{1}{\sqrt{2}}\right)\frac{1}{\gamma}, \quad b = (3 - 2\sqrt{2})\frac{1}{\gamma}, \tag{4.8}$$

and the majorizing function is

$$h(t) = \beta - t + \frac{\gamma t^2}{(1 - \gamma t)}. \tag{4.9}$$

So its two positive solutions are

$$\left. \begin{matrix} r_1 \\ r_2 \end{matrix} \right\} = \frac{1 + \alpha \mp \sqrt{(1 + \alpha)^2 - 8\alpha}}{4\gamma}, \tag{4.10}$$

where $\alpha = \beta\gamma$. From Theorem 3.1 we again have the Smale's type theorem.

Theorem 4.2. Suppose that $r > r_1$, $\|f'(x_0)^{-1}f''(x_0)\| = 2\gamma$ and

$$\begin{aligned} \|f'(x_0)^{-1}(f''(x') - f''(x))\| &\leq \frac{2\gamma}{(1 - \gamma\|x - x_0\| - \gamma\|x' - x\|)^3} - \frac{2\gamma}{(1 - \gamma\|x - x_0\|)^3}, \\ \forall x \in \overline{B(x_0, r)}, \quad \forall x' \in \overline{B(x, r - \|x - x_0\|)}. \end{aligned} \tag{4.11}$$

Let $\beta = \|f'(x_0)^{-1}f(x_0)\|$ and $\alpha = \beta\gamma \leq 3 - 2\sqrt{2}$. Then the iteration

$$x_{n+1} = T_{f,\lambda}(x_n), \quad n = 0, 1, \dots,$$

is well defined for all $\lambda \in [0, 2]$ and $\{x_n\}$ converges to a solution x^* of the equation $f(x) = 0$ satisfying

$$x^* \in \overline{B(x_0 - f'(x_0)^{-1}f(x_0), r_1 - \beta)} \subset \overline{B(x_0, r_1)}.$$

Moreover, for each r satisfying $r_1 \leq r < r_2$ if $\alpha < 3 - 2\sqrt{2}$ and $r = r_1$ if $\alpha = 3 - 2\sqrt{2}$, the equation $f(x) = 0$ has a unique solution in the closed ball $\overline{B(x_0, r)}$.

4.3 Other Examples

Let c be a positive constant. Then, for different functions L given in the following table, we can get a series of concrete results. The corresponding r_0 and b , which have been computed in [17], are illustrated in the following table.

$L(u)$	γr_0	γb
$c\gamma e^{\gamma u}$	$\log \frac{c+1}{c}$	$(c+1) \log \frac{c+1}{c} - 1$
$\frac{c(m+1)\gamma}{(1-\gamma u)^{m+2}}$	$1 - \left(\frac{c}{1+c}\right)^{\frac{1}{m+1}}$	$1 + c \frac{m+1}{m} \left(1 - \left(\frac{c+1}{c}\right)^{\frac{m}{m+1}}\right)$
$\frac{c\gamma}{(1-\gamma u)^2}$	$\frac{1}{c+1}$	$1 - c \log \frac{c+1}{c}$
$\frac{c\gamma}{1-\gamma u}$	$1 - e^{-\frac{1}{c}}$	$1 - c + ce^{-\frac{1}{c}}$

where $m > -1$, $m \neq 0$. Thus using Theorem 3.1 we obtain the corresponding convergence results.

References

- [1] I. K. Argyros and D. Chen, Results on the Chebyshev method in Banach spaces, *Proyecciones*, **12**(1993), 119-128.
- [2] I. K. Argyros, D. Chen and Q. Qian, Optimal-order parameter identification in solving nonlinear systems in a Banach space, *J. Comput. Math.*, **13**(1995), 267-280.
- [3] V. Candela and A. Marquina, Pecurrence relations for rational cubic methods I: The Halley method, *Computing*, **44**(1990), 169-184.
- [4] V. Candela and A. Marquina, Pecurrence relations for rational cubic methods II: The Chebyshev method, *Computing*, **45**(1990), 355-367.
- [5] D. Chen, I. K. Argyros and Q.S. Qian, A note on the Halley method in Banach spaces, *Appl. Math. Comput.*, **58**(1993), 215-224.
- [6] D. Chen, I. K. Argyros and Q.S. Qian, A local convergence theorem for the Super-Halley method in a Banach space, *Appl. Math. Lett.*, **7**(1994), 49-52.
- [7] J. M. Gutierrez and M. A. Hernandez, Convex acceleration of Newton's method, Pub Sem G Galdeano, Serie II, Seccion 1, No 8, Univ. Zaragoza, 1995.
- [8] J. M. Gutierrez and M. A. Hernandez, A family of Chebyshev-Halley type methods in Banach spaces, *Bull. Aust. Math. Soc.*, **55**(1997), 113-130.
- [9] Han Danfu, The convergence on a family of iterations with cubic order, *J. Comput. Math.*, **19**(2001), 467-474.
- [10] M. A. Hernandez, A note on Halley's method, *Numer. Math.*, **59**(1991), 273-276.
- [11] M. A. Hernandez, Newton-Raphson's method and convexity, *Zb Rad Prirod Mat Fak Ser Mat* **22-1**(1992), 159-196.
- [12] M. A. Hernandez and M. A. Salanova, A family of Chebyshev-Halley type methods, *Intern. J. Computer Math.*, **47**(1993), 59-63.
- [13] Wang Xinghua, Convergence of the iteration of Halley's family in weak condition, *Chinese Science Bulletin*, **42**(1997), 552-555.
- [14] Wang Xinghua and Han Danfu, Criterion α and Newton's method in the weak condition, *Math Numer Sinica*, **19**(1997), 103-112; *Chinese J. Numer. Math. Appl.*, **19**(1997), 96-105.
- [15] Wang Xinghua, Convergence of Newton's method and inverse function in Banach spaces, *Math. Comput.*, **68**(1999), 169-186.
- [16] Wang Xinghua, Convergence of Newton's method and uniqueness of the solution of equation in Banach spaces, *IMA J. Numer. Anal.*, **20**(2000), 123-134.
- [17] Wang Xinghua, Convergence of the iteration of Halley family and Smale operator class in Banach space, *Science in China (ser. A)*, **41**(1998), 700-709.
- [18] T. Yamamoto, On the method of tangent hyperbolas in Banach spaces, *J. Comput. Appl. Math.*, **21**(1988), 75-86.